

### **Použitá a doporučená literatura**

- Andersen, L.H., Nummi, P., Rafn, J., Frederiksen, C.M.S., Kristjansen, M.P., Lauridsen, T.L., Trøjelsgaard, K., Pertoldi, C., Bruhn, D., Bahrndorff, S., 2021. Can reed harvest be used as a management strategy for improving invertebrate biomass and diversity? *J. Environ. Manage.* 300, 113637. <https://doi.org/10.1016/j.jenvman.2021.113637>
- Badiou, P.H.J., Goldsborough, L.G., 2015. Ecological impacts of an exotic benthivorous fish, the common carp (*Cyprinus carpio* L.), on water quality, sedimentation, and submerged macrophyte biomass in wetland mesocosms. *Hydrobiologia* 755, 107–121. <https://doi.org/10.1007/s10750-015-2220-6>
- Beckerman, A.P., Wieski, K., Baird, D.J., 2007. Behavioural versus physiological mediation of life history under predation risk. *Oecologia* 152, 335–343. <https://doi.org/10.1007/s00442-006-0642-6>
- Broyer, J., Curtet, L., 2012. Biodiversity and fish farming intensification in French fishpond systems. *Hydrobiologia* 694, 205–218. <https://doi.org/10.1007/s10750-012-1162-5>
- Eichenberg, D., Bowler, D.E., Bonn, A., Bruelheide, H., Grescho, V., Harter, D., Jandt, U., May, R., Winter, M., Jansen, F., 2021. Widespread decline in Central European plant diversity across six decades. *Glob. Chang. Biol.* 27, 1097–1110. <https://doi.org/10.1111/gcb.15447>
- Francová, K., Šumberová, K., Kučerová, A., Čtvrtlíková, M., Šorf, M., Borovec, J., Drozd, B., Janauer, G.A., Vrba, J., 2019. Macrophyte assemblages in fishponds under different fish farming management. *Aquat. Bot.* 159, 103131. <https://doi.org/10.1016/j.aquabot.2019.103131>
- Kajgrová, L., Blabolil, P., Drozd, B., Roy, K., Regenda, J., Šorf, M., Vrba, J., 2022. Negative effects of undesirable fish on common carp production and overall structure and functioning of fishpond ecosystems. *Aquaculture* 549. <https://doi.org/10.1016/j.aquaculture.2021.737811>
- Kłoskowski, J., 2012. Fish stocking creates an ecological trap for an avian predator via effects on prey availability. *Oikos* 121, 1567–1576. <https://doi.org/10.1111/j.1600-0706.2011.19942.x>
- Kłoskowski, J., 2010. Fish farms as amphibian habitats: factors affecting amphibian species richness and community structure at carp ponds in Poland. *Environ. Conserv.* 37, 187–194. <https://doi.org/10.1017/S0376892910000494>
- Kłoskowski, J., Nieoczym, M., Stryjecki, R., 2020. Between-habitat distributions of pond tadpoles and their insect predators in response to fish presence. *Hydrobiologia* 847, 1343–1356. <https://doi.org/10.1007/s10750-020-04190-5>
- Kolář, V., Boukal, D., 2020. Dobré a další zprávy o ochraně ohroženého vodního hmyzu: potápník dvojčárý v ČR. *Fórum Ochr. přírody* 3, 28–32.

- Kolar, V., Boukal, D.S., 2020. Habitat preferences of the endangered diving beetle *Graphoderus bilineatus*: implications for conservation management. Insect Conserv. Divers. 13, 480–494. <https://doi.org/10.1111/icad.12433>
- Kolar, V., Tichanek, F., Tropek, R., 2021a. Evidence-based restoration of freshwater biodiversity after mining: Experience from Central European spoil heaps. J. Appl. Ecol. 1365-2664.13956. <https://doi.org/10.1111/1365-2664.13956>
- Kolar, V., Vlašánek, P., Boukal, D.S., 2021b. The influence of successional stage on local odonate communities in man-made standing waters. Ecol. Eng. 173, 106440. <https://doi.org/10.1016/j.ecoleng.2021.106440>
- Křivánek, J., Němec, J., Kopp, J., 2012. Rybníky v České republice. Pro Ministerstvo zemědělství ČR vydal Consult, Praha.
- Mantefel, Y.B., Reshetnikov, A.N., 2002. Avoidance of noxious tadpole prey by fish and invertebrate predators: Adaptivity of a chemical defence may depend on predator feeding habits. Fundam. Appl. Limnol. 153, 657–668. <https://doi.org/10.1127/archiv-hydrobiol/153/2002/657>
- Mocq, J., Soukup, P.R., Näslund, J., Boukal, D.S., 2021. Disentangling the nonlinear effects of habitat complexity on functional responses. J. Anim. Ecol. 90, 1525–1537. <https://doi.org/10.1111/1365-2656.13473>
- Musil, M., Novotná, K., Potužák, J., Hůda, J., Pechar, L., 2014. Impact of topmouth gudgeon (*Pseudorasbora parva*) on production of common carp (*Cyprinus carpio*) — question of natural food structure. Biologia (Bratisl). 69, 1757–1769. <https://doi.org/10.2478/s11756-014-0483-4>
- Musil, P., Fuchs, R., 1994. Changes in abundance of water birds species in southern Bohemia (Czech Republic) in the last 10 years. Hydrobiologia 279–280, 511–519. <https://doi.org/10.1007/BF00027881>
- Pechar, L., 2000. Impacts of long-term changes in fishery management on the trophic level water quality in Czech fish ponds. Fish. Manag. Ecol. 7, 23–31. <https://doi.org/10.1046/j.1365-2400.2000.00193.x>
- Pykal, J., Janda, J., 1994. Početnost vodních ptáků na jihočeských rybnících ve vztahu k rybničnímu hospodaření. Sylvia 30, 3–11.
- Sánchez-Bayo, F., Wyckhuys, K.A.G., 2019. Worldwide decline of the entomofauna: A review of its drivers. Biol. Conserv. 232, 8–27. <https://doi.org/10.1016/j.biocon.2019.01.020>
- Scheffer, M., Carpenter, S., Foley, J.A., Folke, C., Walker, B., 2001. Catastrophic shifts in ecosystems. Nature 413, 591–596. <https://doi.org/10.1038/35098000>
- Šigutová, H., Šigut, M., Dolný, A., 2018. Phenotypic plasticity in specialists: How long-spined larval *Sympetrum depressiusculum* (Odonata: Libellulidae) responds to combined predator cues. PLoS One 1–15.
- Šigutová, H., Šigut, M., Dolný, A., 2015. Intensive fish ponds as ecological traps for dragonflies: an imminent threat to the endangered species *Sympetrum depressiusculum*

(Odonata: Libellulidae). J. Insect Conserv. 19, 961–974. <https://doi.org/10.1007/s10841-015-9813-2>

Svidenský, R., Kučerová, A., Čížková, H., 2021. Causes of the dieback of littoral stands in an overpopulated water-bird reserve: Role of eutrophication, fish and geese. Eur. J. Environ. Sci. 11, 79–90. <https://doi.org/10.14712/23361964.2021.9>

Tolonen, K.T., Hämäläinen, H., Holopainen, I.J., Mikkonen, K., Karjalainen, J., 2003. Body size and substrate association of littoral insects in relation to vegetation structure. Hydrobiologia 499, 179–190.

Zambrano, L., Hinojosa, D., 1999. Direct and indirect effects of carp (*Cyprinus carpio* L.) on macrophyte and benthic communities in experimental shallow ponds in central Mexico, in: Shallow Lakes '. Springer Netherlands, Dordrecht, pp. 131–138. [https://doi.org/10.1007/978-94-017-2986-4\\_13](https://doi.org/10.1007/978-94-017-2986-4_13)