

Mazuch M., Sakala J.: Tenkrát, když začalo dnešní chladno... (Živa 2021, 5: 212–215)

Použitá a doporučená literatura:

Deutsch A. & Koeberl C. (2006): Establishing the link between the Chesapeake Bay impact structure and the North American tektite strewn field: The Sr-Nd isotopic evidence.

Meteoritics and Planetary Science, Volume41, Issue5, DOI: 0.1111/j.1945-5100.2006.tb00985.x

Gingerich P. & Martin R. (1981): Cranial Morphology and Adaptations in Eocene Adapidae. II. The Cambridge Skull of *Adapis parisiensis*. American Journal of Physical Anthropology, 56: 235-257, DOI: 10.1002/ajpa.1330560304.

Golonka J. (2009): Phanerozoic paleoenvironment and paleolithofacies maps: Cenozoic. Journal Geologia/Akademia Górniczo-Hutnicza im. Stanisława Staszica w Krakowie. 35: 507–587

Hansen J., Sato M., Russell G., Kharecha P. (2013): Climate sensitivity, sea level and atmospheric carbon dioxide. Phil Trans R Soc A 371: 20120294.

DOI: 10.1098/rsta.2012.0294

Houben A.J.P., Bijl P.K., Sluijs A., Schouten S. & Brinkhuis H. (2019). Late Eocene Southern Ocean cooling and invigoration of circulation preconditioned Antarctica for full-scale glaciation. Geochemistry, Geophysics, Geosystems, 20: 2214–2234.

DOI: 10.1029/2019GC008182

Hutchinson D.K., Coxall H.K., Lunt D.J., Steinthorsdottir M., de Boer A.M., Baatsen M., von der Heydt A., Huber M., Kennedy-Asser A.T., Kunzmann L., Ladant J.-B., Lear C.H., Moraweck K., Pearson P.N., Piga E., Pound M.J., Salzmann U., Scher, H.D., Sijp W.P., Śliwińska K.K., Wilson P.A., and Zhang, Z. (2021): The Eocene–Oligocene transition: a review of marine and terrestrial proxy data, models and model–data comparisons. Clim. Past, 17:269–315, DOI: 10.5194/cp-17-269-2021,

Janis C.M. (1993): Tertiary Mammal Evolution in the Context of Changing Climates, Vegetation, and Tectonic Events. Annual Review of Ecology and Systematics. 24: 467–500, DOI: 10.1146/annurev.es.24.110193.002343

Kolektiv autorů (2011): Eocene locality Kučlín. Sborník Národního muzea v Praze, B, 67 (3-4). <https://publikace.nm.cz/periodicke-publikace/fiamnpsbhn/67-3-4>

Kvaček, Z., Walther, H. (2004): Oligocene flora of Bechlejovice. Sborník Národního muzea v Praze, B, 60 (1-2): 9–60.

- Kvaček, Z., Teodoridis, V., Mach, K., Příkryl, T., Dvořák, Z. (2014): Tracing the Eocene–Oligocene transition: a case study from North Bohemia. *Bulletin of Geosciences*, 89(1): 21–66.
- Lefebvre V., Donnadiou Y., Goddérís Y., Fluteau F., Hubert-Théou L. (2013): Was the Antarctic glaciation delayed by a high degassing rate during the early Cenozoic? *Earth and Planetary Science Letters*, 371–372: 203–211, ISSN 0012-821X, DOI: 10.1016/j.epsl.2013.03.049.
- Nelson C. & Cooke P. (2001). History of oceanic front development in the New Zealand sector of the Southern Ocean during the Cenozoic – a synthesis. *New Zealand Journal of Geology and Geophysics*, 44(4): 535-553, DOI: 10.1080/00288306.2001.9514954
- Prothero D. (2003): The Late Eocene-Oligocene Extinctions. *Annual Review of Earth and Planetary Sciences*, 22:145-165. DOI: 10.1146/annurev.earth.22.050194.001045.
- Stehlin H.G. (1910): Remarques sur les faunules de Mammifères des couches eocenes et oligocenes du Bassin de Paris. *Bulletin de la Société Géologique de France*. 4 (9):488–520. (in French).
- Stein R. (2019): The Late Mesozoic-Cenozoic Arctic Ocean Climate and Sea Ice History: A Challenge for Past and Future Scientific Ocean Drilling. *Paleoceanography and Paleoclimatology*, 32(12): 1851-1894, DOI: 10.1029/2018PA003433
- Teodoridis, V., Kvaček, Z. (2015): Palaeoenvironmental evaluation of Cainozoic plant assemblages from the Bohemian Massif (Czech Republic) and adjacent Germany. *Bulletin of Geosciences* 90(3): 695–720.
- Zachos J., Pagani M., Sloan L., Thomas E., Billups K. (2001): Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science* 292:686–693, DOI: 10.1126/science.1059412